



Soils Map



Soils data provided by USDA and NRCS.



State: **Iowa**
 County: **Ringgold**
 Location: **11-67N-31W**
 Township: **Clinton**
 Acres: **72.54**
 Date: **7/28/2026**



MIDWEST
LEGACY
 Land & Farm Sales

Maps Provided By:



© AgriData, Inc. 2025

www.AgriDataInc.com



Area Symbol: IA159, Soil Area Version: 31

Code	Soil Description	Acres	Percent of field	CSR2 Legend	Soil Drainage	Non-Irr Class *c	*i Water-Holding Inch	CSR2**	CSR	*n NCCPI Overall
179E2	Gara loam, 14 to 18 percent slopes, moderately eroded	21.93	30.2%		Well drained	Vle	10.7	24	33	62
792D2	Armstrong clay loam, 9 to 14 percent slopes, moderately eroded	14.68	20.2%		Somewhat poorly drained	IVe	9.0	7	13	58
451D2	Caleb loam, 9 to 14 percent slopes, moderately eroded	5.59	7.7%		Moderately well drained	IVe	9.1	41	33	77
570C	Nira silty clay loam, 5 to 9 percent slopes	4.99	6.9%		Moderately well drained	IIle	11.7	72	69	91
870B	Sharpsburg silty clay loam, terrace, 2 to 5 percent slopes	4.70	6.5%		Moderately well drained	Ile	11.9	91	87	90
179D2	Gara loam, 9 to 14 percent slopes, moderately eroded	3.31	4.6%		Well drained	IVe	10.7	38	43	66
792C2	Armstrong clay loam, 5 to 9 percent slopes, moderately eroded	3.08	4.2%		Somewhat poorly drained	IIle	9.2	24	27	60
76D	Ladoga silt loam, 9 to 14 percent slopes	2.87	4.0%		Moderately well drained	IIle	11.8	52	57	75
76C2	Ladoga silty clay loam, 5 to 9 percent slopes, moderately eroded	2.38	3.3%		Moderately well drained	IIle	11.8	73	62	82
W	Water	2.36	3.3%				0.0	0	0	
452C	Lineville silt loam, 5 to 9 percent slopes	1.57	2.2%		Somewhat poorly drained	IIle	10.8	48	36	75
175D	Dickinson fine sandy loam, 9 to 14 percent slopes	1.39	1.9%		Well drained	IVe	5.3	19	27	57

Soils data provided by USDA and NRCS.

Code	Soil Description	Acres	Percent of field	CSR2 Legend	Soil Drainage	Non-Irr Class *c	*i Water-Holding Inch	CSR2**	CSR	*n NCCPI Overall
179E	Gara loam, 14 to 18 percent slopes	1.03	1.4%		Well drained	Vle	10.7	30	35	66
592D2	Mystic clay loam, 9 to 14 percent slopes, moderately eroded	0.85	1.2%		Somewhat poorly drained	IVe	10.5	10	5	67
570C2	Nira silty clay loam, 5 to 9 percent slopes, moderately eroded	0.68	0.9%		Moderately well drained	IIIe	11.7	68	64	86
13B	Olmitz-Zook-Humeston complex, 0 to 5 percent slopes	0.59	0.8%		Poorly drained	IIw	11.7	78	59	79
269B	Humeston silty clay loam, 2 to 5 percent slopes, rarely flooded	0.54	0.7%		Poorly drained	IIIw	10.1	71	53	88
Weighted Average						*-	10	33.6	36.3	*n 66.3

**IA has updated the CSR values for each county to CSR2.

*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

*n: The aggregation method is "Weighted Average using all components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

*- Non Irr Class weighted average cannot be calculated on the current soils data due to missing data.

	Symbol	Name	Description
	GLA	Glacial till spot	An area of unsorted, nonstratified, glacial drift consisting of clay, silt, sand, cobbles, stones, and boulders. These areas may or may not be calcareous. Typically .25 to 2 acres.
	SAN	Sandy spot	A spot where the surface layer is loamy fine sand or coarser in areas where the surface layer of the named soils in the surrounding map unit is very fine sandy loam or finer. Typically .25 to 2 acres.